

An in vivo photodynamic therapy with diode laser to cell activation of kidney dysfunction

by Suryani Dyah Astuti

Submission date: 08-Nov-2018 03:49PM (UTC+0800)

Submission ID: 1035214259

File name: An_in_vivo_photodynamic_therapy_with_diode_laser_to_cell.pdf (1.01M)

Word count: 3303

Character count: 17105

An in vivo photodynamic therapy with diode laser to cell activation of kidney dysfunction

4

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 J. Phys.: Conf. Ser. 853 012038

(<http://iopscience.iop.org/1742-6596/853/1/012038>)

View the [table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 120.188.67.235

This content was downloaded on 08/06/2017 at 06:53

Please note that [terms and conditions apply](#).

You may also be interested in:

19

[Photodynamic therapy of pigmented melanoma B16 using sterically stabilized fullereneosomes](#)

M Babincová, P Sourivong, D Leszczynska et al.

[Experimental determination of threshold dose in photodynamic therapy in normal rat liver](#)

J Ferreira, L T Moriyama, C Kurachi et al.

10

[A novel frequency domain fluorescence technique for determination of triplet decay times](#)

H J C M Sterenborg, M E Janson and M J C van Gemert

[The dosage optimization of He-Ne laser energy as a candidate for photodynamic therapy of cancer cells with exogenous photosensitizer variations](#)

R Apsari, I Maghfiroh, H Setyawati et al.

[Method for estimating optimal spectral and energy parameters of laser irradiation in photodynamic therapy of biological tissue](#)

S A Lisenko and M M Kugeiko

13

[Light activated antimicrobial agents can inactivate oral malodour causing bacteria](#)

Marika Rai, David Spratt, Paola R Gomez-Pereira et al.

[Tunable 397 nm Light Source for Laser Cooling of Ca Ions Based on Frequency Doubling of Diode Laser](#)

6

Kazuhiro Hayasaka, Masayoshi Watanabe, Hidetsuka Imajo et al.

[Encapsulation of palladium porphyrin photosensitizer in layered metal oxide nanoparticles for photodynamic therapy against skin melanoma](#)

Zih-An Chen, Yaswanth Kuthati, Ranjith Kumar Kankala et al.

An *in vivo* photodynamic therapy with diode laser to cell activation of kidney dysfunction

Suryani Dyah Astuti^{1,2,*}, Brahma Indra Prasaja¹ and Tri Anggono Prijo¹

¹Department of Physics, Faculty of Science and Technology, Airlangga University

²Department of Biomedical Engineering Post Graduate School, Airlangga University

*Corresponding email address: suryanidyah@gmail.com

Abstract. This study aims to analyze the effect of photodynamic therapy (PDT) low level laser therapy (LLLT) 650 nm in the experimental animals mice (*Mus musculus*) suffering from kidney organ damage in mice (*Mus musculus*) in vivo. Exposure laser acupuncture was performed on the kidney BL-23. The conditioning of kidney damage in mice used carbofuran 35 at a dose of 0.041697 mg/mice. LLLT 650 nm exposure was done on a wide variety of energy (0.5; 1.0; 1.5; 2.0; 4.0; 5.0; 6.0; 7.0) J. The histopathological kidney cells in mice renal impairment showed that exposure to 650 nm laser energy 1 Joule resulted in the reduction of damaged cells (necrosis) and normal cells were increased with the improvement of renal tubular cells (64.14 ± 8.02)%. Therefore, exposure to 650 nm LLLT on acupuncture points Shenshu (BL-23) has the ability to proliferation of renal tubular cells of mice.

1. Introduction

Since 2008, biostimulation laser, which is now referred to as Low Level Laser Therapy (LLLT), has been applied to medical therapy [1]. He-Ne laser is the first laser used for wound healing. Biostimulation laser is a photobiology phenomenon that involves chemical reactions and enzymes [2]. Photodynamic activation for biostimulation cells using LLLT is laser stimulation of the skin with low intensity and a non-thermal laser irradiation. Deep transmission of laser energy, wavelength, and laser energy are the important factors on therapy. In additional, scattering and absorption by various tissues of different skins need to be considered. The skin is a network of multi-layer in a homogeneous, anisotropic structure; so that the optical properties are very complex [3]. The intensity of light through the skin decreases exponentially due to the scattering and the absorption. Ultraviolet and visible spectrum (~400-600) nm will be absorbed by hemoglobin and melanin, so, a little amount of the light is transmitted back. For blue light, the irradiance depth would be decreased 99% to ~700 µm. Wavelengths above 1400 nm is the area of water absorption that results on less transmission. Red and infrared spectrum are the most widely used for photodynamic biomodulation because the absorption is very small and the transmission is large [4]. However, scattering by collagen is still considered because it could reduce transmission.

Kidney Disease Outcomes Quality Initiative of The National Kidney Foundation defines that chronic renal failure is a state of the occurrence of kidney damage or glomerular filtration rate (GFR) <60 mL/min/1.73 m² within 3 months or more [5]. The disorder is caused by dysfunction of the



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

glomerular filter or blood and protein filter in the primary urine. Generally, dialysis or hemodialysis could be upgraded to kidney transplant in order to treat chronic kidney failure. Alternative therapy to restore organ function is photodynamic activation through photobiomodulation of kidney cells with a laser exposure. This laser exposure can be performed on acupuncture points, which also called Laserpunctur therapy [6]. Laser therapy acupuncture has many advantages. It is non-invasive, free cause of infection, and environment-friendly. It does not cause pain because of its needle pricks, reduces the risk of disease transferred through a needle, and radiates warm feelings during the therapy. Furthermore, it also causes high immune system response, high growth factor, and is safe for all ages [7]. Based on research, specific acupuncture points used for kidney treatment is shensu (BL-23) [8]. This paper discusses the influence of laser phototherapy at 650 nm in testing animals, which are mice (*Musmusculus*) suffered from kidney disorders, on the kidney acupuncture points BL-23.

2. Materials and methods

Animals used in this study were mice (*Musmusculus*), male, 3-4 week old, and weighing 25-35 grams. *Musmusculus* were conditioned with damaged kidney by giving carbofuran 3% on the dosage of 0.5 mg/kg weight *Musmusculus* [9]. LD₅₀ fraction did not cause the death of the tested animals, yet caused kidney damage is 1/24 LD₅₀ (0.0208 mg / kg), 1/12 LD₅₀ (0.0417 mg/kg) and 1/6 LD₅₀ (0.0833 mg/kg). Therefore, kidney of mice (*Musmusculus*) were set to damaged condition by using Carbofuran 3% with a dose of 0.041697 mg/mice.

2.1. Light sources

Laser irradiation was carried out using diode lasers with output wavelengths of (650.00±0.01) nm. The power outputs were (16.53 ± 0.06) mW. Those data were the result of characterization using Jasco monochromator CT-10 and OPM (Optical Power Meter) Thorlabs PM100D which were capable of detecting power between 100 pW – 200 W and a wavelength of 185 nm – 25 µm. The laser energy was calculated from the output power multiplied by the length of time the laser exposure [11].

2.2. Laser treatment

To determine the laser effect on *Musmusculus*, the samples were distributed into 3 groups, as follows: (1) Groups A was treated using carbofuran and laser 405 nm (C+L+), (2) Groups B was treated only with carbofuran (C+L-), and (3) Group C had no treatment of everything (C-L-). For each group the experiment was repeated at least 3 times. Group A was treated with various energy irradiating of diode laser (1.0; 1.5; 2.0; 4.0; 5.0; 6.0; 7.0) J to acupuncture point kidney (BL-23). Figure 1 shows the acupunctured point BL-23.




Figure 1. Accupuncture point of kidney BL-23

2.3. Histopathological anatomy test

Histopathology test was used to determine the structure of the cells of the renal tubules of mice that had been given laser irradiation treatments as well as healthy and pain control. Analytical calculation of renal tubular cell was conducted using a visual field [10]. The formula to calculate the percentage of regenerated normal cell is by using equation:

$$\% \text{ of normal tubular cells} = \frac{\bar{x} \text{ control diseased cells} - \bar{x} \text{ number of normal cells}}{\bar{x} \text{ control diseased cells}} \times 100\%$$

3. Results and discussion

Laser characterization results in this study are showed in Figure 2 and 3.




Figure 2. Spectrum of 650 nm diode laser





Figure 3. Graph of 650 nm laser power stability versus time

The characterization results in Figure 2 show that the diode laser light source has a spectrum (650.00 ± 0.01) in nm. Figure 3 shows the data output on power stability (16.53 ± 0.06) mW

3.1 Histopathological analysis results

Giving carbofuran orally in mice resulted in infiltration of inflammatory cells and necrosis in renal cells which are indicated by damage within kidney cells [10]. Necrosis is the death of cells in living tissue due to the toxins which enter the body. Inflammation is a complex biological response of vascular tissues to harmful stimuli, such as pathogenic agents, cell death/damaged, or irritation. Inflammation is the body armor attempting to deal with adverse stimuli and to initiate the healing process [12].

In this study, to stimulate the healing process of renal cell, the therapy applied 650 nm laser diode. This therapy was examined through seven variation of energy dose with considerably similar laser beam output because the laser mounted on Shensu acupuncture point and BL-23 had the same cross-sectional area. Figure 4a to 4f illustrate histopathological picture of *Mus musculus*' kidney glomerulus for normal cells, diseased cells, and cells that were exposed to a laser with a magnification of 400X.



Exp: A=glomerular, B=tubulus, C=normal cell, D=Necrosis

Figure 4a. Histopathology normal kidney cells



Exp: A=glomerular, B=tubulus, C=normal cell, D=Necrosis, E=Inflamed cell

Figure 4b. Histopathology disease kidney cells




Exp: A=glomerular, B=tubulus, C=normal cell, D=Necrosis

Figure 4c. Histopathology kidney cells with treatment laser 650 nm 0.5 J




Exp: A=glomerular, B=tubulus, C=normal cell, D=Necrosis

Figure 4d. Histopathology kidney cells with treatment laser 650 nm 1.0 J



Exp: A= glomerulus, B = Tubulus, C = normal cell, D = Necrosis

Figure 4e. Histopathology kidney cells with treatment laser 650 nm 1.5 J



Exp: A= glomerulus, B = Tubulus, C = Normal cell, D = Necrosis

Figure 4f. Histopathology kidney cells with treatment laser 650 nm 2.0 J

Figures 5a and 5b demonstrate the graph of the number of normal cells and damaged at various laser exposure.




Figure 5a. The effect of various energy laser exposure on the number of damaged renal cells in mice

Note:

C+L+ treated with carbofuran and laser 650 nm

C+L- treated with carbofuran and no laser treatment

C-L- no treatment laser and carbofuran




Figure 5b. The effect of various energy laser exposure on the number of normal cells in mice

Figure 5a shows the increase in the number of normal renal cells of mice as a Gaussian function caused by exposure to 650 nm laser for various energy. The laser exposure of 650 nm wavelength and 1.5 J energy dose at acupuncture point BL-23 determines a decrease in the damaged cells the most. As indicated by the Gaussian function, $y = y_0 + \frac{A}{w\sqrt{\pi/2}} e^{-2\frac{(x-x_c)^2}{w^2}}$.

Meanwhile, the number of normal cells in the kidneys of diabetic mice as a result of exposure to 650 nm laser decreases exponentially compared to the number of normal cells if exposed toward the laser with precisely 1.0 J energy. The exponential function is $y = y_0 + A_1 e^{-x/t_1} + A_2 e^{-x/t_2}$ with R^2 value of 0.96.

Figure 6 describes the percentage of renal cell improvement based on histopathological picture.




Figure 6. Growth percentage of renal tubular cells

Figure 6 figures out that the use of a 650 nm laser with 1 Joule energy has improved its percentage ($64.14 \pm 8.02\%$), which is the most optimum result in the improvement of renal tubular cells. The exponential function is $y = y_0 + A_1 e^{-x/t_1} + A_2 e^{-x/t_2}$ with R^2 value of 0.93.

The results of this study showed that a low power laser (LLLT) at a wavelength of 650 nm with 1 J energy at acupuncture points shenshu (BL-23) had the ability to improve renal tubular cells. This is consistent with Peplow's research that shows that LLLT is able to modulate cell proliferation [13, 14]. The results also showed biomodulation effects of LLLT in myofibroblast T and B cells during wound healing. LLLT facilitates differentiation myofibroblastic during the early stages of the cicatricial repair process. Additionally, LLLT also appears to modulate the inflammatory response by down regulating lymphocyte proliferation during the wound healing process [15].

Wavelength laser parameters also affect the ability of cell proliferation. Laser wavelength associated with the depth of penetration into skin. Some research show that wavelength affects on the biological response of HeLa cells, which are irradiated with light of 580-860 nm [16]. Results of other studies also demonstrate the use of laser 780 nm is not as effective as 660 nm laser, yet it has a positive effect on the early stages of the emergence and development of inflammation. At the end of the experimental period, major effect is seen on the amount and quality of granulation tissue. If laser 660 nm is applied daily, it will be more effective than the 780 nm laser to improve third-degree burnt healing in diabetic rats started in the early stages of post-combustion [17-18].

Senshu acupuncture points (BL-23) is a bladder meridian acupuncture points related to the kidney [19]. The bladder is associated with the internal and external kidneys. Stimulation of the points BL-23 provides strengthening effect on the kidneys and kidney bioenergy balance, which in turn will restore kidney function.

4. Conclusion

Based on the results of histopathological kidney cells in mice with renal impairment, it can be concluded that exposure to 650 nm laser with 1 J energy results in a reduction of damaged cells (necrosis) and increase of normal cells with the improvement of renal tubular cells ($64.14 \pm 8.02\%$). Therefore, the exposure to 650 nm LLLT on acupuncture points Shenshu (BL-23) has the ability to proliferation of renal tubular cells of mice.

References

- [1] Karu T I, 1999 Primary and secondary mechanisms of action of visible-to-near IR radiation on cells J. Photochem. Photobiol. B Biol. **49**:1-17.
- [2] Anderson R R 1993 Optics of the skin. In: Lim HW, Soter NA (eds) Clinical photomedicine Marcel Dekker New York pp 19-35
- [3] Kolarova H, Ditrichova D and Wagner J 1999 Penetration of laser light into the skin in vitro *Lasers Surg Med* **24**:231-235 DOI 10.1002/(SICI)1096-9101
- [4] Kidney Disease Outcomes Quality Initiative 2002 Clinical Practice Guidelines For Chronic Kidney Disease: Evaluation, Classification and Stratification New York National Kidney Foundation Inc
- [5] Gerhard L and Gerhard O 2012 Technical Parameters for Laser Acupuncture to Elicit Peripheral and Central Effects: State of the Art and Short Guidelines Based on Result from the Medical University of Graz, the German Academy of Acupuncture, and the Scientific Literature, *Research Article* vol. **2012** Hindawi Publishing Corporation Germany.
- [6] Baxter, G David, C Bleakley and S McDonough 2008 *Clinical Effectiveness of Laser Acupuncture A Systematic Review Journal of Acupuncture and Meridian Studies* **1**(2): 65-82
- [7] He, Tian-Feng, Wen-Jia Yang, Shu-Hui Zhang, Chun-Yan Zhang, Lian-Bo Li and Yun-Fei Chen 2011 Electroacupuncture Inhibits Inflammation Reaction by Upregulating Vasoactive Intestinal Peptide in Rats with Adjuvant-Induced Arthritis Shanghai: Shanghai University of

Traditional Chinese Medicine.

- [8] Kamboj S S, Kumar, A Kamboj and R Sandhir 2008 Mitochondrial Oxidative Stress and Dysfunction in Rat Brain Induced by Carbofuran Exposure *Cell Mol Neurobiol* **28**: 961-969
- [9] Baker J H, Lindsey J K, Steven H W 1979 *The Laboratory Rat: Biology and Disease*. **1**. Unisted States of America: Academic Press.
- [10] Neimz, Markolf H 2007 *Laser-Tissue Interactions Fundamentals and Applications Third, Enlarged Edition*. Germany: Springer
- [11] Kaur, Bhupindervis, Alka Khera and Rajat Sandhir 2012 *Attenuation of Cellular Antioxidant Defense Mechanisms in Kidney of Rats Intoxicated with Carbofuran* *India Journal of Biochemical and Molecular Toxicology*. **26**
- [12] Philip V Peplow, Tzu-Yun Chung, and G David Baxter *Laser Photobiomodulation of Proliferation of Cells in Culture: A Review of Human and Animal Studies* *Photomedicine and Laser Surgery* August 2010, **28**(S1): S-3-S-40. doi:10.1089/pho.2010.2771
- [13] Philip V. Peplow, Tzu-Yun Chung, and G David Baxter, *Laser Photobiomodulation of Wound Healing: A Review of Experimental Studies in Mouse and Rat Animal Models* *Photomedicine and Laser Surgery*. June 2010, **28**(3): 291-325. doi:10.1089/pho.2008.2446.
- [14] Maria Amalia Gonzaga Ribeiro, Ricardo Luiz Cavalcanti Albuquerque, Jr Luciana Maria Pedreira Ramalho, Antonio Luiz Barbosa Pinheiro, Leonardo Rigoldi Bonjardim, and Samantha Seara Da Cunha, *Immunohistochemical Assessment of Myofibroblasts and Lymphoid Cells During Wound Healing in Rats Subjected to Laser Photobiomodulation at 660 nm*, *Photomedicine and Laser Surgery* March 2009, **27**(1): 49-55. doi:10.1089/pho.2007.2215.
- [15] T I Karu, S F Kolyakov, *Exact Action Spectra for Cellular Responses Relevant to Phototherapy*, *Photomedicine and Laser Surgery* 2005, **23**(4): 355-361
- [16] G C S Meireles, J N Santos, P O Chagas, A P Moura, A L B Pinheiro, *Effectiveness of Laser Photobiomodulation at 660 or 780 Nanometers on the Repair of Third-Degree Burns in Diabetic Rats* *Photomedicine and Laser Surgery* 2008 **26**(1):47-54 DOI: 10.1089/pho.2007.2051
- [17] Brodie, L M 2003 *Welding of Skin using Nd:YAG Laser with Bipolar Contact Applicators*. *Queensland* thesis of University of Southern Queensland Faculty of Engineering and Surveying
- [18] Z Yanfu, *Chinese Accupuncture and Moxibustion*, Shanghai University of Traditional Chinese Medicine 2002

An in vivo photodynamic therapy with diode laser to cell activation of kidney dysfunction

ORIGINALITY REPORT

23%

SIMILARITY INDEX

20%

INTERNET SOURCES

18%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	pubman.mpdl.mpg.de Internet Source	4%
2	www.er-c.org Internet Source	2%
3	www.laser-therapy.us Internet Source	2%
4	www.lps.ens.fr Internet Source	2%
5	www.healinglightseminars.com Internet Source	2%
6	Mei Yu, Zhiqin Yuan, Chao Lu. "Layered double hydroxide-enhanced luminescence in a Fenton-like system for selective sensing of cobalt in Hela cells", Methods and Applications in Fluorescence, 2017 Publication	1%
7	blog.thorlaser.com Internet Source	1%

8

D V Vysotskii, N N Elkin, A P Napartovich,
Vladimir I Kozlovskii, B M Lavrushin. "Modelling
of a diode laser with a resonant grating of
quantum wells and an external mirror",
Quantum Electronics, 2011

Publication

1 %

9

S A Lisenko, M M Kugeiko, V A Firago, A N
Sobchuk. "Analytical model of diffuse
reflectance spectrum of skin tissue", Quantum
Electronics, 2014

Publication

1 %

10

H J C M Sterenborg. Physics in Medicine and
Biology, 05/1999

Publication

1 %

11

V V Kardumyan, N A Aksenova, A A Chernyak,
N N Glagolev, V I Volkov, A B Solovieva. "The
influence of temperature on the photo-
oxidation rate of tryptophan in the presence of
complexes of porphyrins with amphiphilic
polymers", Laser Physics, 2015

Publication

1 %

12

sylvain-delzon.com

Internet Source

1 %

13

Subhangi Mittal, Sukhdev Roy, J N Srivastava.
" Fungicidal response of a novel natural
photosensitizer () on with low-power laser
radiation ", Laser Physics, 2013

1 %

14

Clementina López-Medina, Yolanda Jiménez-Gómez, Anna Moltó, Ruxandra Elena Schiotis et al. "Cardiovascular risk factors in patients with spondyloarthritis from Northern European and Mediterranean countries: An ancillary study of the ASAS-COMOSPA project", Joint Bone Spine, 2017

Publication

1 %

15

Chan, Hsiu-Hui, Tsong-Long Hwang, Tran Thang, Yann-Lii Leu, Ping-Chung Kuo, Bui Nguyet, Do Dai, and Tian-Shung Wu. "Isolation and Synthesis of Melodamide A, a New Anti-inflammatory Phenolic Amide from the Leaves of Melodorum fruticosum", Planta Medica, 2013.

Publication

1 %

16

isan.troitsk.ru

Internet Source

<1 %

17

ar.orthodoxebooks.org

Internet Source

<1 %

18

www.biomedcentral.com

Internet Source

<1 %

19

M. Babinová. "Photodynamic therapy of pigmented melanoma B16 using sterically stabilized fullerenosomes", Laser Physics

<1 %

-
- | | | |
|---|---|----------------|
| <div style="background-color: #003366; color: white; display: inline-block; width: 40px; height: 40px; text-align: center; line-height: 40px;">20</div> | pure.iiasa.ac.at
<small>Internet Source</small> | <1 % |
|---|---|----------------|
-
- | | | |
|---|---|----------------|
| <div style="background-color: #800080; color: white; display: inline-block; width: 40px; height: 40px; text-align: center; line-height: 40px;">21</div> | anatomydatabase.otago.ac.nz
<small>Internet Source</small> | <1 % |
|---|---|----------------|
-
- | | | |
|---|--|----------------|
| <div style="background-color: #6B8E23; color: white; display: inline-block; width: 40px; height: 40px; text-align: center; line-height: 40px;">22</div> | Metin Güngörmüş. "Effect of Biostimulation on Wound Healing in Diabetic Rats",
Photomedicine and Laser Surgery, 08/2009
<small>Publication</small> | <1 % |
|---|--|----------------|
-
- | | | |
|---|---|----------------|
| <div style="background-color: #000080; color: white; display: inline-block; width: 40px; height: 40px; text-align: center; line-height: 40px;">23</div> | Rinaldo Roberto de Jesus Guirro. "Analysis of Low-Level Laser Radiation Transmission in Occlusive Dressings", Photomedicine and Laser Surgery, 08/2010
<small>Publication</small> | <1 % |
|---|---|----------------|
-
- | | | |
|---|---|----------------|
| <div style="background-color: #0056b3; color: white; display: inline-block; width: 40px; height: 40px; text-align: center; line-height: 40px;">24</div> | worldwidescience.org
<small>Internet Source</small> | <1 % |
|---|---|----------------|
-
- | | | |
|---|---|----------------|
| <div style="background-color: #ff0000; color: white; display: inline-block; width: 40px; height: 40px; text-align: center; line-height: 40px;">25</div> | www.isan.troitsk.ru
<small>Internet Source</small> | <1 % |
|---|---|----------------|
-
- | | | |
|---|---|----------------|
| <div style="background-color: #cc00cc; color: white; display: inline-block; width: 40px; height: 40px; text-align: center; line-height: 40px;">26</div> | elib.uni-stuttgart.de
<small>Internet Source</small> | <1 % |
|---|---|----------------|
-
- | | | |
|---|---|----------------|
| <div style="background-color: #6a0dad; color: white; display: inline-block; width: 40px; height: 40px; text-align: center; line-height: 40px;">27</div> | hub.hku.hk
<small>Internet Source</small> | <1 % |
|---|---|----------------|
-
- | | | |
|---|---|----------------|
| <div style="background-color: #008080; color: white; display: inline-block; width: 40px; height: 40px; text-align: center; line-height: 40px;">28</div> | Gyselle C.S. Meireles. "Effectiveness of Laser Photobiomodulation at 660 or 780 Nanometers | <1 % |
|---|---|----------------|

on the Repair of Third-Degree Burns in Diabetic Rats", Photomedicine and Laser Surgery, 02/2008

Publication

29

Maria Amalia Gonzaga Ribeiro.
"Immunohistochemical Assessment of Myofibroblasts and Lymphoid Cells During Wound Healing in Rats Subjected to Laser Photobiomodulation at 660 nm",
Photomedicine and Laser Surgery, 02/2009

<1%

Publication

30

G. Arun Maiya. "Effect of Low Intensity Helium-Neon (He-Ne) Laser Irradiation on Diabetic Wound Healing Dynamics", Photomedicine and Laser Surgery, 04/2005

<1%

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

An in vivo photodynamic therapy with diode laser to cell activation of kidney dysfunction

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8